

Lucile Packard Children's Hospital Stanford

Lucile Salter Packard Children's Hospital



Page 1 of 2

Patient Name: Date of Birth:

Well Baby Check: 12 month visit questionnaire

Interval History: Has your child had any major illnesses or doctor visits since last seen	n here? No	Yes		
Has your child had any reactions to vaccinations in the past?		Yes		
Development : Can your child (check all that apply) -				
_ hold a cup to drink?	_ look at something when you point and say "look"?			
_ feed him/herself finger foods?s	_ see well without crossing or drifting eyes?			
_ pick objects up with thumb and index finger? _ w	walk while holding onto furniture (cruise)?walk alone?clap hands?			
_ combine syllables (e.g. "dada," "mama")?				
_ use gestures (point with finger/hand)? _ c				
_ understand words ("no," "more")?	_ wave bye-bye?			
_ play peek-a-boo?				
Who provides daytime care for your child?				
Nutrition/Elimination:				
How much milk does your child drink? oz per day of [breast m	ilk] [formula] [v	whole mil	k] [other	
Does your baby get 3 servings of calcium-rich foods daily?	Yes	No		
s your child eating a variety of fruits and vegetables daily?	Yes	No		
Does your child eat iron-rich foods (meat/iron-fortified cereal, beans) daily? Yes	No		
Do you give your baby a bottle of anything except formula, milk or	water? No	Yes		
Do you offer your child a sippy cup every day?	Yes	No		
Are there any problems with pooping or peeing?	No	Yes		
Baby's medications/vitamins/supplements:				
Mother's medications/vitamins/supplements if giving breastmilk:				
Dental Health:				
Do you brush your child's teeth daily?	Yes	No		
Sleep:				
How long does your baby sleep at night without awakening?		_ hours		
How long does your baby nap during the day?		_ hours		
Does you baby sleep through the night without feeding?	Yes	No		
Can they self-soothe?	Yes	No		
Where does your baby sleep?				
Staying Healthy/Safety:				
Does your baby get any screen time?	No	Yes		
Does your home have a working smoke detector?	Yes	No		
Is your water temperature set to less than 120 degrees?	Yes	No	N/A	
Is your haby always supervised when near water, including the batht	uh? Vec	No		

L15859 (11/20)



Lucile Packard Children's Hospital

Stanford

Lucile Salter Packard Children's Hospital

STANFORD UNIVERSITY MEDICAL CENTER • 725 Welch Road, Palo Alto, CA 94304

Questionnaire • Well Baby Check 12 Month Page 2 of 2 Patient Name:

Date of Birth:

Have you child-proofed your home?	Yes	No	
Do you have safety guards on upper floor windows and gates for the stairs?	Yes	No	N/A
Does your home have cleaning supplies/medicines/matches locked away?	Yes	No	
Is the Poison Control Center number (800-222-1222) posted by/in your phone?	Yes	No	
Does your child use sun protection when outdoors?	Yes	No	
Is your car seat appropriately sized, rear-facing, and in the back seat?	Yes Yes	No No	
Are all guns stored in a gun safe or locked with ammunition separate from gun?			N/A
Does your baby spend time with anyone who smokes or vapes?	No	Yes	
Risk Assessment for Lead Exposure:			
Does your child live or spend a lot of time in a place built before before 1978			
that has peeling or chipped paint?	No	Yes	
Does your child live or spend a lot of time in a place built before before 1978			
that has been recently renovated?	No	Yes	
Please list any new major family medical issues:			
Who lives in the home with your child?			
What international travel has your child had since their last well check? (v	where a	and how	long)
What plans are there for international travel with your child in the next 12	month	us? (whe	ere and how long)
What concerns would you like to discuss today?			

(11/20)L15859



Lucile Packard Children's Hospital Stanford

Lucile Salter Packard Children's Hospital STANFORD UNIVERSITY MEDICAL CENTER • 725 Welch Road, Palo Alto, CA 94304



Outpatient Record • TB Skin Test

Patient Name

Patient Date of Birth

School:					
Grade: Santa Clara	County	Public Health Departm	ent		
Tuberculosis (TB)	Risk As	sessment for School En	try		
This form must be completed by a licensed he	-		rned to th	e child's scho	
 Was your child born in, resided, or traveled (country with an elevated rate of TB*? 	your child born in, resided, or traveled (for more than one month) to a with an elevated rate of TB*?			□ No	
2. Has your child been in close contact to anyoutheir lifetime?	as your child been in close contact to anyone with tuberculosis (TB) disease in lifetime?			□ No	
3. Is your child immunosuppressed; current, or organ transplant, treatment with TNF-alpha ant steroids (e.g. prednisone ≥ 15 mg/day for ≥ 2 w *Most countries other than the U.S., Canada, At This does not include tourist travel for <1 month significant contact with the local population).	agonist or reeks). ustralia, N	high-dose systemic ew Zealand, or a country in we			
If YES, to any of the above questions, the child i.e. QuantiFERON or T-SPOT.TB) or a tuberculir IGRA or TST performed in the U.S. or 2) no new ≥2 years in the U.S.) or TST (performed at age ≥	n skin test risk facto	(TST) unless there is either 1) rs since last documented nega	a docume	nted prior positi	
All children with a current or prior positive IGRA/(CXR; posterior-anterior and lateral for children < documented prior treatment for TB disease, documented prior treatment for TB disease, documented who have a positive TST and negative Idnormal, the child should be treated for latent TB in	5 years our	ld is recommended). CXR is no rior treatment for latent TB infe ere are no symptoms or signs o	ot required ection, or B of TB disea	for children with BCG-vaccinated ase and the CXI	
Enter test results for all children with a positi	ve risk as	sessment:			
Interferon Gamma Release Assay (IGRA)					
Date:		Result: Negative Pos	sitive 🛭 I	ndeterminate	
Tuberculin Skin Test (TST/Mantoux/PPD) Induration mm					
Date placed: Date read:		Result: Negative Pos	sitive		
Chest X-Ray Date: Impression	n: 🗖 Norr	nal 🛘 Abnormal			
LTBI Treatment Start Date:		☐ Prior TB/LTBI treatment (Rx & duration):			
 □ Rifampin daily - 4 months □ Isoniazid/Rifapentine - weekly X 12 weeks □ Isoniazid daily - 9 months □ Other: 		☐ Treatment medically contraindicated			
		☐ Declined against medical advice			
		Declined against medical	auvice		
Please check one of the boxes below and sign: Child has no TB symptoms, no risk factors Child has a risk factor, has been evaluated Child has no new risk factors since last neg	for TB, and	d is free of active TB disease.			
	Health C	Care Provider Signature, Title	<u>-</u>	Date	
Name/Title of Health Provider:		 			
License Number: Facility/Address:					
Phone number:					

County of Santa Clara

Public Health Department

Tuberculosis Prevention & Control Program 976 Lenzen Avenue, Suite 1700 San José, CA 95126 408.885.2440



Testing Methods

An Interferon Gamma Release Assay (IGRA, i.e., QuantiFERON or T-SPOT.TB) or Mantoux tuberculin skin test (TST) should be used to test those at increased risk. An IGRA can be used in all children \geq 2 years old and is preferred in BCG-vaccinated children to avoid a false positive TST result. A TST of \geq 10mm induration is considered positive. If a child has had contact with someone with active TB disease (yes to question 2 on reverse), or the child is immunosuppressed, then TST \geq 5 mm is considered positive. If a BCG-vaccinated child has a positive TST, and an IGRA is subsequently performed and is negative, testing is considered negative unless the child was exposed to someone with TB disease or is immunosuppressed. For immunosuppressed children, screening should be performed by CXR in addition to a TST/IGRA (consider doing both) and symptom review. TB screening can be falsely negative within 8 weeks after exposure, so are best obtained 8 weeks after last exposure.

Evaluation of Children with Positive TB Tests

- All children with a positive IGRA/TST result must have a medical evaluation, including a CXR (posterior-anterior
 and lateral is recommended for children <5 years old). A CXR is not required for a positive TST with negative
 IGRA in a BCG-vaccinated child, or if the child has documentation of prior treatment for TB disease or treatment
 for latent TB infection.
- For children with TB symptoms (e.g., cough for >2-3 weeks, shortness of breath, hemoptysis, fever, weight loss, night sweats) or an abnormal CXR consistent with active TB disease, report to the County of Santa Clara Public Health Department TB Program within one day. The child will need to be evaluated for TB disease with sputum AFB smears/cultures and nucleic acid amplification testing. A negative TST or IGRA does not rule out active TB disease in a patient with symptoms or signs of TB disease. The child cannot enter school unless active TB disease has been excluded or treatment has been initiated.
- If there are no symptoms or signs of TB disease and the CXR is normal, the child should be treated for latent TB infection (LTBI). Do not treat for LTBI until active TB disease has been excluded.
- Short-course regimens (rifampin daily for four months or 12-dose weekly isoniazid/rifapentine) are preferred (except in persons for whom there is a contraindication, such as a drug interaction or contact to a person with drug-resistant TB) due to similar efficacy and higher treatment completion rates as compared with 9 months of daily isoniazid.

Treatment Regimens for Latent TB Infection

- Rifampin 15 20 mg/kg (max. 600 mg) daily for 4 months
- 12-dose Weekly Isoniazid/Rifapentine (3HP) Regimen:
 - Isoniazid

2-11 years old: 25 mg/kg rounded up to nearest 50 or 100 mg (max. 900 mg) ≥ 12 years old: 15 mg/kg rounded up to nearest 50 or 100 mg (max. 900 mg)

Rifapentine

10.0-14.0 kg: 300 mg 14.1-25.0 kg: 450 mg 25.1-32.0 kg: 600 mg 32.1-50.0 kg: 750 mg >50 kg: 900 mg

Vitamin B6 50 mg weekly

 Isoniazid 10 mg/kg (range, 10-15 mg/kg; max. 300 mg) daily for 9 months. Recommended pyridoxine dosage is 25 mg for school-aged children (or 1-2 mg/kg/day).

Board of Supervisors: Mike Wasserman, Cindy Chavez, Otto Lee, Susan Ellenberg, S. Joseph Simitian, County Executive: Jeffrey V. Smith